

# Spiny water flea (*Bythotrephes longimanus*)

**Species and Origin:** Spiny waterfleas are zooplankton (microscopic animals) that are native to Europe and Asia.

**Date of U.S Introduction:** They were first discovered in Lake Ontario in 1982. They were introduced into the Great Lakes by ballast water discharged from ocean-going ships.

**Date of Minnesota Introduction:** They spread to Lake Superior in 1987.

**Description:** Adults range from 1/4 to 5/8 inch long. Spiny waterfleas have a single long tail with small spines along its length.

**Impacts:** Spiny waterfleas eat small animals (zooplankton), including *Daphnia*, which are an important food for native fishes. In some lakes, they caused the decline or elimination of some species of native zooplankton. They can clog eyelets of fishing rods and prevent fish from being landed.

**Status:** They have spread throughout the Great Lakes, and are established in some inland lakes and rivers in Minnesota.

**Means of spread:** They can spread by attaching to fishing lines, downriggers, anchor ropes, and fishing nets. While female waterfleas die out of water, under certain conditions they produce eggs that resist drying and freezing, and can establish a new infestation. They also can be unintentionally transported in bilge water, bait buckets, or livewells.

**Where to look:** They collect in gelatinous globs on fishing lines and downrigger cables. They prefer deep lakes, but can be found in shallow lakes and rivers.



## *Bythotrephes longimanus*

Map created on 11/8/2011. United States Geological Survey

